

NOAA Technical Report NMFS Circular 412



Synopsis of Biological Data  
on the Red Porgy,  
*Pagrus pagrus* (Linnaeus)

May 1978



FAO Fisheries  
Synopsis No. 116  
NMFS/S 116  
SAST - Red Porgy  
1.70(39) 191.03

U.S. DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service



NOAA Technical Report NMFS Circular 412  
**Synopsis of Biological Data  
on the Red Porgy,  
*Pagrus pagrus* (Linnaeus)**

Charles S. Manooch III and William W. Hassler

May 1978

FAO Fisheries Synopsis No. 116

**U.S. DEPARTMENT OF COMMERCE**

Juanita M. Kreps, Secretary

**National Oceanic and Atmospheric Administration**

Richard A. Frank, Administrator

**National Marine Fisheries Service**

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

# CONTENTS

|   | Page |
|---|------|
| Introduction . . . . .  | 1    |
| 1 Identity . . . . .  | 1    |
| 1.1 Nomenclature . . . . .                                      | 1    |
| 1.2 Taxonomy . . . . .  | 1    |
| 1.3 Morphology . . . . .  | 3    |
| 2 Distribution . . . . .  | 3    |
| 2.1 Total area . . . . .  | 3    |
| 2.2 Differential distribution . . . . .                         | 4    |
| 2.3 Determinants of distribution . . . . .                      | 5    |
| 3 Bionomics and life history . . . . .                          | 5    |
| 3.1 Reproduction . . . . .                                      | 5    |
| 3.2 Preadult phase . . . . .                                    | 6    |
| 3.3 Adult phase . . . . .                                       | 7    |
| 3.4 Nutrition and growth . . . . .                              | 9    |
| 3.5 Behavior . . . . .  | 10   |
| 4 Population . . . . .  | 10   |
| 4.1 Structure . . . . .   | 10   |
| 4.2 Abundance and density . . . . .                             | 11   |
| 4.4 Mortality and morbidity . . . . .                           | 12   |
| 4.6 The population in the community and the ecosystem . . . . . | 13   |
| 5 Fishing . . . . .   | 13   |
| 5.1 Fishing equipment . . . . .                                 | 13   |
| 5.2 Fishing areas . . . . .                                     | 14   |
| 5.3 Fishing seasons . . . . .                                   | 16   |
| 5.4 Fishing operations and results . . . . .                    | 17   |
| Acknowledgments . . . . .                                       | 17   |
| Literature cited . . . . .                                      | 18   |

# Synopsis of Biological Data on the Red Porgy, *Pagrus pagrus* (Linnaeus)

CHARLES S. MANOOCH III<sup>1</sup> and WILLIAM W. HASSLER<sup>2</sup>

## ABSTRACT

A synopsis of the biology of the red porgy, *Pagrus pagrus*, that includes taxonomy, morphology, distribution, aspects of the life history, behavior, and abundance. Also included are: discussions of commercial and recreational fishing methods and fishing grounds, and size, age, and sex composition of the recreational catch off North Carolina and South Carolina and commercial catch off Argentina.

## INTRODUCTION

In recent years, the Fisheries Biology Branch of FAO has formed a "Synopsis Association" composed of various fishery agencies which have assisted in preparation of synopses on aquatic organisms of economic importance. Some of these synopses have been published in the U.S. Fish and Wildlife Service and the National Marine Fisheries Service Circular series following the format prescribed by Rosa (1965).

The major purpose of the synopses is to make existing information on various species of fishes and invertebrates easily accessible to scientists and the fishery constituencies according to a standard format. Gaps in knowledge will be exposed, thus delineating areas of research needs.

### 1 IDENTITY

#### 1.1 Nomenclature

##### 1.11 Valid name

*Pagrus pagrus* (Linnaeus 1758)

##### 1.12 Synonymy

- Sparus pagrus* Linnaeus 1758:279, southern Europe.  
*Sparus orphus* Linnaeus 1758:279, no locality.  
*Sparus argenteus* Bloch and Schneider 1801:27, South America.  
*Pagrus orphus*. Valenciennes 1837-1844:32, Canary Islands.  
*Pagrus pagrus*. Valenciennes 1837-1844:32, Canary Islands.  
*Pagrus pagrus*. Jordan and Gunn 1899:341, Canary Islands.

*Pagrus sedecim* Ginsburg 1952:93, Pensacola, Fla.

#### 1.2 Taxonomy

##### 1.21 Affinities

|          |                      |
|----------|----------------------|
| Phylum   | Chordata             |
| Class    | Osteichthyes         |
| Order    | Perciformes          |
| Suborder | Percoidae            |
| Family   | Sparidae             |
| Genus    | <i>Pagrus</i>        |
| Species  | <i>Pagrus pagrus</i> |

#### Generic

The generic description follows Fowler (1936): "Body oblong, compressed. Head large. Eye small. Mouth rather small, terminal, low. Front teeth in jaws cardiform, outer series usually enlarged, canine-like, not compressed; teeth behind canines slender and acute; both jaws with 2 or 3 series of rounded molar teeth, sometimes irregularly mixed with slender teeth; no teeth on vomer or palatines. Hind nostril oblong, not slitlike, much larger than anterior. Preorbital deep. Preopercle entire. Opercle not armed. Gill rakers short. Branchiostegals 6. Air vessel simple. Intestinal canal short. Pyloric appendages short. Scales moderately large. Dorsal rather low, spines 11 or 12, depressible in groove; rays 10 or 11. Anal spines 3, moderate, rays 8. Caudal forked."

Another important generic character is: "... dorsal fin not preceded by a projecting spine (present in *Calamus*, but largely hidden by skin and scales)..." (Randall 1968).

Four valid species of *Pagrus* are presently recognized, two from the eastern Atlantic *P. auriga* and *P. ehrenbergi*, one from both eastern and western Atlantic, *P. pagrus* (Fig. 1), and one species from the Indo-Pacific (*P. major*). Briefly, the geographical range for each species is as follows:

*Pagrus pagrus* (Linnaeus)—Eastern Atlantic from the

<sup>1</sup>Southeast Fisheries Center Beaufort Laboratory, National Marine Fisheries Service, NOAA, Beaufort, NC 28516.

<sup>2</sup>Department of Zoology, North Carolina State University, Raleigh, NC 27607.




Figure 1.—Red porgy, *Pagrus pagrus*, approximately 425 mm total length caught off North Carolina (drawing by Herberg Gordy, NMFS).

British Isles, Azores and Canary Islands south to Angola; western Atlantic from New York south to Argentina; includes the Gulf of Mexico but not the West Indies or Central America.

*Pagrus auriga* (Valenciennes)—Eastern Atlantic from Madeira and the Canary Islands to Guinea.

*Pagrus ehrenbergi* (Valenciennes)—Eastern Atlantic and Mediterranean, south to Angola.

*Pagrus major* (Temminck and Schlegel)—Yellow Sea, Sea of Japan; on the Pacific coast of Japan from Hokkaido southward; East and South China seas southward to Australia and in the Indian Ocean.

### Specific

Dorsal fin rays 12-13 (usually 12), 9-11 (usually 10); anal fin rays 3, 7-9 (usually 8); lateral-line scales 51-59; total first-arch gill rakers 13-17, usually  $6 + 1 + 9 = 16$ ; pectoral rays 14-16; vertebrae  $10 + 14 = 24$ ; branchiostegal rays 6. Body moderately deep, depth 2.59 to 2.93 times in standard length (SL); pectoral fin 2.59 to 3.50 times in SL; head 2.76 to 3.60 times in SL; eye diameter 2.77 to 5.48 times in head; nostril width 1.05 to 4.25 in nostril length; caudal peduncle depth 8.11 to 10.92% SL; maxillary with cleft midlaterally; length of head 0.8428 to 0.8821 in head depth; head length 0.2585 to 0.2829 in distance from the base of the orbit to the jaw articulation (depth of cheek). Meristics and proportional measurements are from Manooch et al. 1976.

Both eastern and western Atlantic specimens are colored similarly, although the heads of eastern Atlantic porgy are darker, particularly after preservation. Overall body color in life is reddish above and silver-white below. When stressed, fish become barred with 5 to 6 red, vertical bands. There are 11 to 12 rows of small blue spots occurring from the nape region to the caudal peduncle and extending ventrally to just below the base of the pectoral fin. Ventrally, the fish are white to gray-white.

There are generally two light blue laterally directed streaks, one just above, and one just below the eye. The nostril and maxillary areas are silver-blue (metallic). The eye consists of a yellow iris and a black pupil. The ventral fins in life are light blue, but fade to white-gray upon death. The pectoral fins are light yellow. The caudal fin is also light yellow shading to red at its posterior edge. The spiny dorsal fin is pink and the soft dorsal is yellow. Underwater, western Atlantic specimens appear uniformly silver-white with the blue spots evident. A dark band runs from just below the eye to the articulation of the jaw. Preserved specimens appear dull, yellow-brown. The blue, red, and distinct yellow colors are no longer evident on the fins. The spots are still visible, but are black or brown (modified from Manooch et al. 1976).

### 1.22 Taxonomic status

Ginsburg (1952) concluded that the form found in the western Atlantic was a separate species, *Pagrus sedecim*, from that in the eastern Atlantic on the basis of number of pectoral rays, relative depth of the caudal peduncle, and presence of blue spots. Ginsburg's small sample sizes and examination of eastern and western Atlantic specimens in the U.S. National Museum of Natural History prompted Manooch et al. (1976) to review the status of *Pagrus sedecim* Ginsburg. Specimens of eastern Atlantic *Pagrus pagrus* were collected from the Canary Islands, Spain, and those of nominal *P. sedecim* were collected off North Carolina and in the Gulf of Mexico. Meristics, proportional measurements, and electrophoresis were used to evaluate whether *P. sedecim* was a separate species. Results indicated intraspecific differentiation; therefore, *P. sedecim* was placed in the synonymy of *P. pagrus*, and eastern and western Atlantic porgy were considered to represent populations of a



single species. Cotrina and Cousseau<sup>3</sup> examined 150 specimens from Mar del Plata, Argentina, and found no significant difference in meristics or morphometrics compared with eastern Atlantic (European) fish. Also, Travassos (1973), working with red porgy from Brazil, found no basis for the species division of eastern and western Atlantic fish as proposed by Ginsburg (1952).

### 1.23 Standard common names, vernacular names.

United States—Red porgy, Pink porgy, Silver snapper, Rose perch  
 Canary Islands—Bocinegro, Bocanegra, Besugo  
 Madeira—Pargo  
 Senegal—Debesjhal  
 Mauritania, Spanish Sahara—Lauriana  
 France—Pagré  
 British Isles—Sea bream  
 Italy—Pagri, Pargo  
 Poland—Pagrus, Rodzaj pagrusy  
 Brazil—Pargo róseo  
 Uruguay—Pargo colorado, Besugo  
 Argentina—Besugo  
 Spain—Pargo  
 Libya—Mamrya  
 Malta—Pagru  
 Tunisia—Hamraia, Gerali  
 Turkey—Mercan  
 Portugal—Pargo  
 Israel—Shefarnun  
 Yugoslavia—Pager  
 Greece—Fagri

## 1.3 Morphology

### 1.31 External morphology

Manooch et al. (1976) derived a series of proportional body measurements and meristic counts from approximately 80 specimens from the Canary Islands, Gulf of Mexico, and North Carolina (Table 1). These data made it possible to compare the meristics and morphometrics of fish from three different geographical areas.

### 1.33 Protein specificity

Manooch et al. (1976) used two types of electrophoretic procedures—regular disc gel and sodium dodecyl sulfate—to try to find differences among 11 species of sparids. Those analyzed were: *Pagrus pagrus*, nominal *P. sedecim*, *P. auriga*, *Pagellus acarne*, *P. erythrinus*, *P. mormyrus*, *Dentex filiosus*, *D. macrophthalmus*, *Diplodus sargus*, *Calamus leucosteus*, and *C.*

Table 1.—Mean meristic and proportional measurements of 11 characters tested by discriminant function analysis differentiating between three areas of collection (from Manooch et al. 1976).

| Character tested             | Collection site     |                     |                     |
|------------------------------|---------------------|---------------------|---------------------|
|                              | North Carolina (25) | Gulf of Mexico (13) | Canary Islands (42) |
| Dorsal rays                  | 9.96                | 9.92                | 9.98                |
| Anal rays                    | 8.00                | 8.08                | 8.02                |
| Pectoral rays                | 15.84               | 15.92               | 15.21               |
| Lateral-line scales          | 54.92               | 54.61               | 53.88               |
| Gill rakers                  | 15.60               | 15.61               | 15.93               |
| SL/depth                     | 2.70                | 2.64                | 2.67                |
| SL/pectoral length           | 2.90                | 2.79                | 2.79                |
| SL/head                      | 3.15                | 3.01                | 3.15                |
| Head length/eye diameter     | 3.95                | 3.68                | 3.67                |
| Nostril length/nostril width | 2.48                | 2.34                | 2.60                |
| Caudal peduncle depth/SL     | 9.14                | 9.92                | 9.66                |

*nodosus*. By using both electrophoretic techniques they found distinct differences among all pairs except *P. pagrus* and *P. sedecim*.

## 2 DISTRIBUTION

### 2.1 Total area

Red porgy is a sublittoral, demersal, marine species which occurs to deep zones of continental shelves. The species is found in the eastern Atlantic from the British Isles (occasionally) south to Angola off the west coast of Africa, in the Azores and Canary Islands, and in the Mediterranean and Adriatic (Hilgendorf 1888; Fowler 1936; Soljan 1963; Ranzi 1969). Red porgy also occur in the western Atlantic and Gulf of Mexico ranging as far north as New York (occasionally) south to Argentina (Randall 1968); the species has not been reported from the Caribbean or Central America (Fig. 2).

Red porgy are commonly found at depths ranging from 18 to 183 m. Off northwest Africa, Murray and Hjort (1912) reported fishermen capturing *Pagrus pagrus* in waters ranging from 33 to 55 m. Exploratory trawling by the Poles produced red porgy in depths ranging from 22 to 150 m (Wozniak 1967; Klimaj 1970). The deepest collection site recorded is 280 m, south of the Canary Islands (Murray and Hjort 1912). Bottom water temperatures around the Canary Islands and on the West African Continental Shelf range from 15° to 20°C annually (Jones and Folkard 1970). Wheeler (1969) described the habitat of *P. pagrus* in the Mediterranean and Atlantic off North Africa: "A sea bream most commonly found on gravel bottoms or around rocks, in depths up to 33 fathoms (60 m), but young specimens can be caught near the shore within the normal range of the species."

In the western Atlantic, red porgy commonly occur over very irregular and low profile, hard bottom. Austin<sup>4</sup> reported that the red porgy was the most commonly

<sup>3</sup>Cotrina, C. P., and M. B. Cousseau. Undated. Informe sobre el muestreo bioestadístico do desembarque de pescado en el Puerto de Mar del Plata, Argentina. Periodo Enero de 1972-Diciembre de 1974. Instituto de Biología Marina, Mar del Plata, Argentina. Unpagued.

<sup>4</sup>Austin, H. M. 1971. Ecology of fishes on Florida's Middle Ground. Fla. State Univ., Dep. Oceanogr., Tallahassee, Fla., 56 p.




Figure 2.—Distribution of *Pagrus pagrus*.

caught species on the Florida Middle Ground, an area in the Gulf of Mexico off Tampa. The habitat is characterized by a flat, sandy bottom in 40 m of water with limestone outcroppings rising 15 m off the bottom. Bottom water temperatures ranged from 13° to 25°C. Manooch (1975) described the habitat of red porgy off North Carolina and South Carolina. Red porgy were caught by hook and line over several types of habitat, e.g., open-shelf, live-bottom, shelf-edge, and lower shelf, but were concentrated over inshore, live-bottom areas, and at the shelf-edge where the bottom is characterized by a series of troughs, terraces, and ridges which parallel the shelf dropoff (Macintyre and Milliman 1970). The inshore concentrations occurred from 24 to 42 m, and offshore from 51 to 128 m. Inshore, red porgy and related species are usually found associated with any irregularity: shipwrecks, rock outcroppings, and small patches of coral (Huntsman and Macintyre 1971), on the otherwise smooth, sandy bottom live-bottom as defined by Struhaker (1969). Bottom temperatures, including both inshore and offshore, range from approximately 14° to 25°C annually (Manooch 1975).

In South American waters red porgy are also found over rough bottom in depths similar to the North American distribution. N. A. Menezes (pers. commun., Museu de Zoologia da Universidade de São Paulo) reported the species occurring from 19 to 144 m off Brazil and Uruguay, and Cotrina and Cousseau (see footnote 3) stated that *P. pagrus* seldom occur deeper than 37 m off Mar del Plata, Argentina. M. Yesaki (pers. commun., SUDEPE/PDP, Rio Grande) found red porgy concentrated in the mixing area of the cold Falkland Current and the warm Brazil Current (13°-18°C) off Brazil during 1973, and S. S. Cordeiro and A. Miyares (pers. com-

mun., Programa de Pesquisa ex Desenvolvimento Pesqueiro do Brasil) stated that *Pagrus* commonly occurred at 40 to 59 m off the southern coast of Brazil.

## 2.2 Differential distribution

### 2.2.1 Eggs, larvae, and juveniles

Studies of early life stages of the red porgy have been hampered by the difficulty of identification at the species level; this has been particularly true in North America. Manooch (1976) provided a basic description of the spawned, unfertilized egg for fish collected off North Carolina. Descriptions of eggs, larvae, and prejuveniles are provided by Ranzi (1969) and Ciechomski and Weiss (1973) for eastern Atlantic and South American stocks, respectively. Since the eggs and larvae are pelagic for a number of days, their distribution is greatly influenced by currents and winds.

Ranzi (1969) referred to vertical migration of larval and postlarval phases of *Pagrus pagrus*: "All of these stages [<10 mm] can be fished in the deep plankton but at 10 mm *Pagrus* comes to the surface." Also he noted a shift from planktonic to benthic existence at lengths above 20 mm.

It is highly probable that young *Pagrus* are distributed inshore of adult populations. Manooch (1975) reported young-of-year trawled from 9 m off Charleston, S.C. Even though they may occur inshore as juveniles, adverse conditions of low water temperature, competitive exclusion, and unsuitable substrate reduce the longevity of this inshore distribution.

Not only are eggs and larvae transported inshore by Ekman transport, but they are probably transported for



relatively long distances. Manooch et al. (1976) proposed a hypothesis explaining population similarities of eastern Atlantic and western Atlantic *Pagrus pagrus*: "... although the populations have been separated a long time sufficient gene flow has occurred to prevent divergence. Passive transport of larvae from Africa is considered a probable mechanism for maintaining gene flow between the eastern and western Atlantic stocks."

## 2.22 Adults

Adult red porgy occupy a wide horizontal and vertical range (section 2.1), preferring hard substrate, both low and high profiles, and do not display marked seasonal movements. Adults do not occur in waters as shallow as larvae and juveniles but the inshore occurrence of young is probably short-lived (section 2.21). Adults do not occur at great depths under the upper layers which apparently transport eggs and larvae.

## 2.3 Determinants of distribution

Substrate: hard bottom; low and high profile. Depth: 9 to 280 m. Temperature: 13° to 26° C.

# 3 BIONOMICS AND LIFE HISTORY

## 3.1 Reproduction

### 3.11 Sexuality

Both protandrous and protogynous hermaphroditism are relatively common among the sparids (D'Ancona 1950). D. S. Beaumariage (pers. commun., Florida Department of Natural Resources) reported red porgy collected off the west coast of Florida appear to display protogynous hermaphroditism, although data at present are insufficient for quantitative description. Manooch (1976) reported that a predominance of females at smaller size intervals (<400 mm total length (TL)), a large proportion of males for the larger size intervals (>450 mm TL), and discovery of individuals with both testicular and ovarian tissue supports the theory of protogyny. Functional hermaphroditic fish were collected off the Carolinas.

### 3.12 Maturity

Manooch (1976) determined age at sexual maturity for female red porgy. The correlation of age with maturity suggests that none of the age I fish, 37% of the age II fish, 81% of age III fish, and 100% of the age IV fish were mature.

### 3.13 Fertilization

Fertilization is external.

## 3.14 Gonads (ovaries)

Manooch (1976) evaluated three predictors of fecundity from 50 fish: total length, weight, and age. All three variables could be used to predict fecundity, but weight of fish proved to be the best predictor. The equation describing the relationship and coefficient of determination is:

$$\ln \text{ fecundity} = 1.7369 + 1.5178 (\ln \text{ weight}), r^2 = 0.70.$$

The 95% confidence limits are also presented (Fig. 3). Point estimates of fecundity ranged from 48,660 eggs for a fish 304 mm TL and weighing 390 g, to 488,600 ova for a female measuring 516 mm TL and weighing 1,783 g. Theoretically, a 600-mm red porgy could produce approximately 943,000 eggs if maximum ova production is not obtained at a smaller size (age).




Figure 3.—Relationship between fecundity and fish weight for 50 red porgy collected from Onslow Bay, N.C. (from Manooch 1976).

## 3.15 Spawning

In their study of egg and larval development, Ciechomski and Weiss (1973) referred to red porgy spawning in the Argentine Sea mainly from December through January when the water temperature range approximated 20° to 21°C. Ranzi (1969) mentioned the period of sexual maturity of *P. pagrus* off Algiers as April to June. Manooch (1976) collected ripe females from Raleigh and Onslow Bays, N.C., over irregular bottom from January through April in water ranging in depth from 21 to 100 m. Bottom temperatures varied from 16.4° to 21.5°C. Peak spawning occurred from March through April. Gonad condition was believed to be more closely associated with photoperiod than bottom water temperature (Fig. 4). Early maturing and ripe stages of testes were easily discernible, but the late maturing and ripe classes were difficult to separate. Milt could be pressed from the central canal of testes from January through April.




Figure 4.—Mean monthly gonad indices for female *Pagrus pagrus* for each month compared with photoperiod and bottom temperature (from Manooch 1976).

### 3.16 Eggs

Ripe, unfertilized red porgy eggs were pelagic, spherical, without appendages, measured 0.64 to 0.92 mm in diameter, and contained a single oil droplet which measured 0.20 to 0.32 mm in diameter (Manooch 1976). Ciechomski and Weiss (1973) described fertilized eggs which measured 0.81 to 0.88 mm in diameter with an oil droplet which measured 0.18 to 0.21 mm in diameter.

### 3.2 Preadult phase

#### 3.21 Embryonic phase

A description of embryonic development is translated from Ciechomski and Weiss (1973) from the Argentine Sea (Fig. 5): The perivitelline space was small and emphasized at the two poles. Two hours after fertilization 16 to 32 blastomeres were observed in the animal pole. The oil droplet was positioned close to the vegetal pole and aided with flotation as the eggs floated close to the water surface. After 5 h a blastula with medium cells formed;




Figure 5.—Larval stages of red porgy from fertilized egg to 4 days after hatching (from Ciechomski and Weiss 1973).

at 8 to 8½ h gastrulation began. In this stage the germinal ring was evident; the embryo was similar to other species of teleosts and started to increase in size. Morphometric changes were rapid and the otic capsules, first myomeres, and brainfolds could be noted. After 17 to 18 h, the embryo occupied the first half of the egg diameter

and its posterior parts began to grow independently from the yolk sac. The first pigmentary cells appeared as small black points distributed along the body and yolk sac. After 21 h, the embryo occupied about two-thirds of the egg's diameter and the optic capsules were formed. The number of myomeres and pigment cells increased, especially in the zone close to the optic capsules and in the caudal region. Twenty-six hours after fertilization, the embryo occupied two-thirds of the perimeter of the egg and the otoliths were visible in the otic capsule. Hatching occurred from 28 to 38 h after fertilization (temperature, 21.5° to 22.5°C).

### 3.22 Larval phase

The following is a description of larvae from hatching to 4 days (Ciechomski and Weiss 1973). The recently hatched larvae measured 2.42 mm and were almost transparent. The yolk sac was large and oval shaped. In most larvae the oil drop was situated in the posterior portion of the yolk sac, but in a few it was found in the center. The anus was located behind the yolk sac in the middle of the body. The digestive tract opened at the anus at a right angle. The fin fold originated behind the head and was continuous to the anus. All capsules, otic, optic, and olfactory, were evident. The same distribution of pigment for all larvae of the same stage was typical for the species. The yellow cells were distributed along the body in five bands. At 1 day the larvae were 3.1 mm, and at 2 days 3.25 mm. When the larvae were 2½ days old the mouth was functional.

Ranzi (1969) described larval and juvenile phases of *P. pagrus* from the Bay of Naples. Specimens collected were pelagic until 20 mm and then shifted to the bottom. Brief descriptions for fish 3.2 to 15 mm follow.

3.2 mm—The tail was a little more than twice as long as the body; the maximum height was contained ca. 3.5 times in the length. General shape of the body was rather slim. The unpaired fin was continuous and the caudal-most area revealed some visible outlines of rays, not perceived in the area of the anal and dorsal. Of the paired fins, only the pectorals were present. On the operculum, a series of spines, typical of later stages, was evident represented by protruding crests. There was no occipital crest. The eye was pigmented, the intestine enlarged anteriorly, bent ventrally, terminated in the anus, and opened at the level of the 10th and 11th myomere.

4.1 mm—The specimen had features of small *Pagrus*; that is, the tail became larger than in the previous stage with respect to the body, accounting for almost two-thirds of its length, while the head became very powerful; the maximum depth was thus contained only 3 times in the length. The anus had moved in the rostral direction and was located at the level of the seventh myomere. On the operculum two series of spines had developed, the posterior more developed with a very long central spine located at the angle where the whole series bent in the dorsal direction. Dorsally to the eye there was a crest with two slight indentations, and a preliminary outline of spines. On the dorsal profile, just behind the head, there

was a strong unpaired spine, typical of young *Pagrus* from this stage up to the length of 15 mm; the belly, caudally to the operculum, had another strong unpaired spine.

8 mm—The tail was half as long as the body. The supraorbital crest had four small teeth; the inner opercular crest had seven short spines and the outer crest eight spines, the fifth the longest. The ventral fins were formed but there were no outlines of rays; the unpaired fins and the pectorals had definitive rays as in the adult: D 12/10; A 3/8; C 17; P 15. The anal was prolonged in front of the rays by a membranous portion and reached the anus.

9 mm—This specimen was similar to the one at 8 mm. All the previous stages could be fished in the deep plankton but at 10 mm *Pagrus* came to the surface.

11 mm—The body color was pinkish-yellowish. Its dorsoventral diameter increased less than the length and was 2.2 mm in the cephalic region, which accounted for one-fifth of the length. In the scapular region a series of short spines appeared. The definitive rays were evident in the ventrals.

13 mm—Transverse stripes appeared. The first eight rays of the dorsal, all the rays of the ventrals, and the three spiny rays of the anal were yellow. Scales appeared on the body.

15 mm—The lateral line of the trunk, convex upward fairly near the dorsal profile, was evident. The supraorbital crest was less visible than in the previous stages. The whole body was covered with scales. Dorsally and just behind the head, the occipital spine was still present, which up to this stage is typical of small *Pagrus* (Fig. 6).

## 3.3 Adult phase

### 3.31 Longevity

Red porgy are slow growing and have a relatively long life span. Although the species may attain an age of 15 yr or older (Manooch and Huntsman 1977; Cotrina and Cousseau see footnote 3), apparently rapid total annual mortality occurs after age XI off the southeastern United States. From 1972 through 1974, NMFS personnel examined over 13,000 red porgy landed by recreational fishermen fishing from North Carolina and South Carolina headboats. Fish aged IV to VIII were common in the catch (Manooch 1975) (Fig. 7).

### 3.32 Hardiness

No quantitative data are available on the tolerance of red porgy to marked changes in the environment. Personal experiences with the species allow us to conclude that *P. pagrus* is perhaps the hardiest of the demersal fishes which inhabit the outer continental shelf off North Carolina and South Carolina. Red porgy have been taken from depths of >100 m, tagged, and then recaptured later with no obvious ill effects caused by pressure difference. In holding tanks, red porgy have undergone rather dras-




Figure 6.—Young *Pagrus pagrus* from 3.2 to 15 mm (from Ramzi 1969).




Figure 7.—Age and length distributions for 13,120 red porgy sampled from the Carolina headboat fishery (from Manooch 1975).

tic changes in temperature, over relatively short periods of time. Changes from 8.0° to 29.1°C have not proven fatal (R. O. Parker, pers. commun., National Marine Fisheries Service, Beaufort, N.C.).

### 3.33 Competitors

Any fish which feeds on similar foods and occupies the same habitat with the red porgy could be considered a competitor. Considering the wide distribution of *P. pagrus*, the list of competitive fish could of course, be very extensive. Off northwest Africa, other sparids could be major competitors: *Dentex*, *Pagellus*, *Diplodus*, *Sparus*, and *Pagrus* are some of the genera. In North Carolina and South Carolina, species of serranids, lutjanids, pomadasyids, and branchiostegids are major competitors for food and, to an extent, space.

### 3.34 Predators

All large carnivorous sea mammals, turtles, and fishes which occur with red porgy are potential predators. Red porgy remains have been identified in stomachs of sharks, barracuda, and greater amberjack (personal observation).



### 3.35 Parasites, disease, injuries, and abnormalities

There is little published information on parasites and diseases of red porgy. Sproston (1946) listed three species of monogenetic trematodes found on the gills of *P. pagrus*: *Anoplodiscus richiardii* (from the Mediterranean), *Diplectanum echineis* (from Trieste), and *Microcotyle archosargi* (from North America, Atlantic). Causey (1953) identified the copepod *Lernanthropus brevoortiae* Rathbun on a fish thought to be *Pagrus se-decim* from the Texas coast.

## 3.4 Nutrition and growth

### 3.41 Feeding

The red porgy is an opportunistic browser which feeds on a tremendous variety of invertebrates as well as small fishes. Manooch (1975) described some aspects of red porgy feeding: "They are fast enough to compete successfully with other fish for such motile foods as small fish, portunid crabs, squids, and shrimp. Also, *P. pagrus* possesses strong molariform teeth which enable it to crush less motile, armored forms such as echinoderms, pagurid crabs, and gastropods. The red porgy, then, seems well adapted to feed on motile forms and also on non-motile organisms which must depend on armour as a protection against predation. Both observations made in large aquaria, and experimental fishing reveal the red porgy to be very aggressive feeders. They generally seize the food (or bait) immediately when it is presented to them." Manooch (in press) also noted that the red porgy is a bottom feeder: "Approximately 74% of the foods were classified as obligate benthic organisms; 60% by volume. If semibenthic organisms are included, the percentages are increased to 90 to 92 respectively."

### 3.42 Food

As mentioned above (section 3.41) red porgy feed on a variety of invertebrates and small fishes. Wheeler (1969) reported that the diet of *Pagrus pagrus* collected in the eastern Atlantic "... consists principally of fish and crustaceans, including crabs, shrimps and prawns. On broken ground large specimens eat quantities of octopus." In the western Atlantic, Bearden and McKenzie (1969) mentioned decapods (Majidae, Xanthidae, and Portunidae), polychaetes, and mollusks in the diet of fish collected off South Carolina, and Austin (see footnote 4) found crabs (*Mithrax pleuracanthus* and *Stenorynchus seticornis*), ophiuroids, and mollusks in stomachs of red porgy collected from the Gulf of Mexico. Red porgy collected from Argentine waters fed on fishes, mollusks, crabs, sessile coelenterates, polychaetes, amphipods, and echinoderms (Cotrina and Cousseau see footnote 3). Manooch (in press) found 69 different food items in red porgy stomachs collected off North Carolina and South Carolina including 14 species of fishes and 55 taxa of in-

vertebrates. Invertebrates, represented primarily by crustaceans, mollusks, and echinoderms, occurred in 89% of the stomachs. Crabs, mainly majids, portunids, and calappids, were the predominant food by both frequency of occurrence and volume. Little difference in the diet was noted between seasons, area of collection, or depth. The diet of juveniles (46-64 mm) varied significantly from adults: these smaller fish consumed amphipods, copepods, stomatopods, isopods, and annelids.

### 3.43 Growth rate

Age and growth has been determined for red porgy collected from North Carolina and South Carolina waters (Manooch and Huntsman 1977). Average lengths for 1,777 red porgy aged by scales collected in 1972 to 1974 were: 238, 290, 341, 382, 419, 451, 483, 505, 527, 543, 558, and 604 mm for ages I to XII, respectively. Back-calculated lengths of 1,622 fish for ages I to XIII were: 179, 259, 316, 358, 389, 417, 440, 451, 476, 491, 509, 534, and 559 mm, respectively. The oldest fish, 15 yr, was 694 mm TL. Growth is more rapid the first 7 yr (Fig. 8). Observed and back-calculated length-age data were used to calculate theoretical growth. Age groups I to XII were used to fit Walford (1946) lines and the von Bertalanffy (1938) growth equation. This equation for Carolina red porgy is:

$$l_t = 746 (1 - e^{-0.096(t + 1.88)}).$$

Manooch and Huntsman (1977) also derived a length-weight relationship for Carolina red porgy. This equation is:

$$W = 0.00002524L^{2.8939}$$




Figure 8.—Absolute growth curves for *Pagrus pagrus* from mean total lengths of each age group obtained by reading scales and otoliths. The number by each point on the curves represents the number of fish sampled (from Manooch 1975).



where  $W$  = weight in grams and  
 $L$  = total length.

The equation  $W = 0.000027836L^{2.8831}$  describes the length relationship for males, and  $W = 0.000013378L^{2.9954}$  is the equation for females. Dias et al. (1972) furnished length-weight relationships for the species off South Carolina, and Cotrina and Cousseau (see footnote 3) provided the following length-weight regression for 1,044 specimens collected from Mar del Plata, Argentina:

$$W = 0.00002056L^{2.897}$$

### 3.5 Behavior

#### 3.51 Migration and local movements

Apparently, red porgy do not undergo long-range migrations, and local movements are not extensive. Trawling off the West African Continental Shelf (Wozniak 1967; Klimaj 1970) identified concentrations of *Pagrus* by depth and geographic area, but it could not be determined if the species migrates from one location to another. Tagging studies off the Carolinas (Manooch 1975) revealed that *P. pagrus* did not move far from the original tagging site. The average distance moved over a 2-yr period was 5.9 km and the farthest a tagged fish moved was 23.7 km after 47 days. Red porgy tagged and returned off of the west coast of Florida revealed no movement (Beaumariage 1969).

#### 3.52 Schooling

Red porgy do occur in schools, however no quantitative data is available on school characteristics.

## 4 POPULATION

### 4.1 Structure

Since 1972, NMFS has obtained information on the sex, age, and size composition of the red porgy population off the Carolinas by regularly sampling the recreational landings. The procedures of the sampling program and catch statistics have been reported (Sekavec and Huntsman 1972; Manooch 1975; Huntsman 1976). The Instituto de Biologia Marina, Mar del Plata, Argentina, has obtained similar population characteristics by sampling commercial landings (Cotrina and Cousseau see footnote 3).

#### 4.11 Sex ratio

The ratio of male to female red porgy landed off North Carolina and South Carolina was not 1:1 but actually 1:2.1. Of the total 705,372 *P. pagrus* landed by the headboat fishery, approximately 259,638 were males and 445,733 were female. The percentage of males to females varied with size of fish and also by season (month). The

greatest discrepancy in the ratio occurred in the spring, and was approximately 1:3. Red porgy spawn in the spring off the Carolinas, and the discrepancy between sexes caught could indicate schooling of females prior to spawning, but it is more probably due to the fact that the species displays protogynous hermaphroditism and more females are available (Manooch 1976). Commercial Argentine landings also reveal a predominance of females in the catch (Cotrina and Cousseau see footnote 3).

#### 4.12 Age composition

Red porgy aged from the recreational fishery off North Carolina and South Carolina ranged from 1 to 15 yr. To facilitate analysis of data, fish were identified in age groups I to XII separately, and those few individuals XIII to XV were categorized as >XII (Manooch and Huntsman 1977).

A histogram of age frequency for 1972 to 1974 indicates that the greatest percentage of red porgy caught by the recreational fishery are age V (Fig. 7). For all years, 29.6% of the porgy landed were 5-yr-old fish. Gear selectivity, rather than relative abundance of any age group, has a definite effect on age composition of the catch. This is particularly true with hook and line where smaller individuals are completely or selectively excluded from the catch.

Age V was the most frequently landed age group each year: 1972, 32.4%; 1973, 28.7%; and 1974, 27.4%. A larger percentage of younger fish (I-IV) were landed in 1974 than in the other years. Whereas, in 1972 and 1973 ages VI and VII composed approximately 28% of the catch, in 1974 ages III and IV were more frequently caught than ages VI and VII.

Stratification of age frequency data by area reveals some differences in age composition. While age V occurred most frequently in each area, the Cape Fear, N.C., catch was composed of a larger percentage of younger fish. Twice as many age I and II red porgy were landed in the Cape Fear, N.C., vicinity as in the Cape Lookout, N.C., and Cape Romain, S.C., districts. Age IV was the next most frequently encountered age group in Cape Fear, whereas age group VI was encountered more frequently in the Cape Lookout and Cape Romain districts. As with size, the occurrence of younger fish in the Cape Fear district is probably correlated with inshore fishing intensity.

Young red porgy are more frequently caught inshore (Fig. 9). Age groups I to IV composed approximately 51.4% of the catch inshore compared with only 24.9% offshore. Hook size varied between the two grounds, and undoubtedly contributed to age composition differences of the catch. However, experimental fishing with a variety of hook sizes (1/0-9/0) aboard the research vessel indicated a predominance of larger (older) fish in deeper waters.

Red porgy aged by scales from Argentina ranged from 0 to 17 yr (Cotrina and Cousseau see footnote 3). Percentages by age for more than 6,500 fish from commercial trap and trawl landings, 1972 to 1974 were: age




Figure 9.—Length-frequency distribution for 13,120 red porgy collected on inshore and offshore fishing grounds off the Carolinas (from Manooch 1975).

II—8.5, III—17.2, IV—22.2, V—11.6, VI—9.4, VII—11.9, VIII—8.2, IX—4.7, X—1.9, and >X—3.9.

#### 4.13 Size composition

Red porgy caught by hook and line off the Carolinas from 1972 to 1974 ranged from 212 to 741 mm (8.3 to 29.2 in) TL. Over half (56.3%) of the fish landed were between 400 and 499 mm (15.7 and 19.6 in) TL and averaged 425.4 mm (16.7 in) (Fig. 7). The length-weight relationship predicts an average weight of 1.0 kg (2.25 lb) for a 425-mm fish.

Size of red porgy varied only slightly between years. Average fish size was largest in 1973; 439.5 mm (17.3 in) TL and 1.1 kg (2.48 lb). Average size for the other years was very similar. The largest red porgy sampled was 741 mm (29.2 in) and 5.1 kg (11.2 lb) landed in South Carolina in 1974.

Mean sizes were again similar by area. Fish averaged slightly larger in the Cape Romain district: 426.7 mm (16.8 in) and 1.0 kg (2.26 lb) compared with 425.0 mm (16.7 in) (0.8 kg (1.83 lb)) and 421.8 mm (16.7 in) (1.0 kg (2.19 lb)) for Cape Lookout and Cape Fear districts, respectively. A different length-weight relationship was used for each area resulting in different mean weights per length of fish. A larger percentage of smaller fish (<350 mm) were caught in the Cape Fear district, and relatively more individuals greater than 600 mm (23.6 in) were landed in South Carolina. One explanation of the occurrence of smaller fish in the Cape Fear vicinity is that a greater fraction of the effort is inshore, where smaller hooks are used and smaller fish are found.

A more pronounced difference in size is revealed by stratifying data according to inshore and offshore fishing grounds. Inshore fish were consistently smaller for each year compared with red porgy caught on the offshore grounds (Fig. 9). Mean size for inshore fish for all years was 387.6 mm (15.2 in) and 0.78 kg (1.72 lb) compared with 436.6 mm (17.2 in) and 1.1 kg (2.43 lb) offshore. Ap-

proximately 69% of the fish collected in shallower water ranged from 325 to 449 mm (12.8-17.7 in), and 68% of the offshore individuals ranged from 400 to 524 mm (15.7-20.6 in) TL.

*Pagrus pagrus* exploited commercially off South America are generally smaller than those sampled from the Carolina recreational fishery. Fish from Argentina ranged from 100 to 540 mm TL, and most were between 260 and 380 mm (Cotrina and Cousseau see footnote 3). Cordeiro and Miyares (pers. commun.) reported that specimens caught commercially by hook and line off Brazil are generally 320 to 380 mm in length.

#### 4.2 Abundance and density

Data on standing crop or biomass per unit area for red porgy are limited. While no information is available from North America or the eastern Atlantic, Yesaki (pers. commun.) gives estimates between 78,000 to 104,000 t in 1973 and 10,000 to 55,000 t in 1974 during December off the southern coast of Brazil.

The NMFS has conducted surveys which report catch and catch per unit effort (CPUE) for *P. pagrus* off the Carolinas. The following discussion is from Manooch (1975): "A recreational fishery exists for the species throughout much of its U.S. range. Most of the effort is from headboat<sup>4</sup> anglers although private and charter vessels also land *Pagrus*. Unfortunately, catch and effort statistics are lacking except for North Carolina and South Carolina, where in 1972 the National Marine Fisheries Service initiated studies of offshore demersal stocks. In 1972, 1973, and 1974 over 700,000 fish weighing 771 metric tons were caught by North Carolina and South Carolina anglers (Sekavec and Huntsman 1972; Huntsman 1976). The species contributed 50% by number, and 44.6% of the weight of all offshore demersal species harvested by the fishery excluding black sea bass, *Centropristis striata*."

Red porgy are also landed commercially; however, the species is generally combined with two other genera of porgy, *Stenotomus* and *Calamus*, under the heading "scup" or "porgy" (Anonymous 1971). The species probably represents only a small percentage of this category, but would be impossible to separate by reviewing these catch records.

Catch and effort data for the years 1972 to 1974 are presented in Table 2 stratified by inshore and offshore fishing. For the 3 yr surveyed, 705,372 red porgy weighing 786.95 t were caught during 193,564 angler days of fishing. This results in a mean catch of 3.64 red porgy and 4.07 kg per angler day.

In 1972, 48,991 angler days were expended to catch 215,958 red porgy which weighed 235.3 t. The average fisherman offshore caught approximately 6 fish (7.26 kg) per angler compared with 2.5 (2.27 kg) fish for anglers fishing inshore.

<sup>4</sup>Headboats are boats where anglers pay for a day's fishing on a per person basis.

Table 2.—Catch and effort data for red porgy stratified by trip type and year, 1972-74 (from Manooch 1975).

| Year | Trip type | Angler days | Number caught | Total weight  |            |            |
|------|-----------|-------------|---------------|---------------|------------|------------|
|      |           |             |               | (metric tons) | No./angler | Wt./angler |
| 1972 | Offshore  | 25,082      | 156,463       | 185           | 6.24       | 16.23      |
|      | Inshore   | 23,909      | 59,495        | 51            | 2.49       | 4.68       |
|      | Subtotal  | 48,991      | 215,958       | 235           | 4.41       | 10.59      |
| 1973 | Offshore  | 42,731      | 227,018       | 280           | 5.31       | 14.45      |
|      | Inshore   | 16,235      | 64,411        | 53            | 3.97       | 7.20       |
|      | Subtotal  | 58,966      | 291,429       | 333           | 4.94       | 12.45      |
| 1974 | Offshore  | 30,867      | 117,828       | 142           | 3.82       | 10.13      |
|      | Inshore   | 54,740      | 80,157        | 77            | 1.46       | 3.09       |
|      | Subtotal  | 85,607      | 197,985       | 219           | 2.31       | 5.64       |
|      | Totals    | 193,564     | 705,372       | 787           | 3.64       | 8.96       |

Fishing was slightly better in 1973. Approximately 59,000 angler days resulted in a catch of 291,429 red porgy which totaled 333.0 t. The average catch per angler day was 4.9 fish weighing 5.65 kg. Again, as in 1972, offshore angling produced a higher CPUE than did inshore fishing: 5.31 fish (6.35 kg) per angler day compared with 4.00 (3.18 kg).

Catch per angler day dropped rather drastically in 1974. Approximately 85,600 angler days resulted in a catch of 197,985 fish which weighed 218.6 t. The average catch of 2.31 fish and 2.55 kg per angler day was the lowest of the 3 yr surveyed. As with the preceding years, fishing was better offshore, 3.82 red porgy per angler day, compared with inshore, 1.46 per angler day. One explanation for the increase in effort and a factor which reduced CPUE is that in 1974, nine additional inshore boats in South Carolina were included in the survey. Not only did these boats increase the effort by 39,248 angler days, but they also fished very shallow areas which are typically black sea bass fishing grounds. Excluding these boats, 160,254 red porgy were harvested by 46,359 angler days; a catch rate of 3.46 fish per angler day.

Catch per unit effort for red porgy increased southward. For all years combined, fishing grounds in the Cape Fear district produced the highest catch per angler day, 4.73 fish and 5.01 kg. Fishing effort was greatest in the Cape Romain district and was actually composed of 47% of the total angler days for all areas combined. Catch per unit effort was also relatively high in South Carolina: 4.14 red porgy and 4.60 kg per angler day. The South Carolina catch per angler day is 6.54 fish and 7.40 kg per angler day if the 1974 inshore boats are excluded. Fishing at Cape Lookout and Cape Hatteras, N.C., was less productive. Only 1.91 fish per angler day were caught in the Cape Lookout district and 2.13 at Cape Hatteras.

Fishing offshore was almost always more rewarding than fishing inshore. Catch per unit effort was better inshore at Cape Hatteras, but there was a small number of angler days fished.

The highest catch rate obtained was 9.08 red porgy and 10.31 kg per angler day, offshore in the Cape Romain area in 1972; and the lowest was 0.66 fish and 0.55 kg per angler day for 1974 anglers, inshore at Cape Lookout.

Catch per unit effort not only varied with year and geographical location, but also between seasons.

Seasonal differences were very inconsistent. For all years and areas combined, spring revealed the highest CPUE, 3.90 red porgy and 4.17 kg per angler day. However, inspection of the combined data reveals little difference in the catch rates: spring, 3.90 porgy and 4.17 kg per angler day; June, 3.84 and 4.07 kg; July, 3.50 and 4.09 kg; August, 3.33 and 3.95 kg; and fall, 3.81 fish and 4.13 kg.

Inspection of these data by year indicates the highest CPUE in the fall for 1972 and 1974 and in the spring for 1973.

Looking at each stratum independently, the highest CPUE occurred offshore in June 1972, 7.47 fish and 8.41 kg per angler day. Unmeasured variables such as weather, currents, and angler quality undoubtedly have pronounced effects on catch per angler day.

#### 4.4 Mortality and morbidity

Total annual mortality rates were calculated for red porgy off North Carolina and South Carolina (Manooch and Huntsman 1977). Age composition of 1,777 red porgy obtained from 1972 to 1974 from the North Carolina and South Carolina headboat fishery was derived by scale reading. The fish were grouped into 25-mm size intervals and the percentage (probability value  $z$ ) for each age (I-XII) was identified for each size interval. Probability values for each age and size interval could then be applied to the length-frequency data derived from 13,120 fish sampled from the sport fishery for the years 1972 to 1974. These data were then stratified by year and area of collection.

Mean total annual mortality estimates for red porgy in North Carolina and South Carolina ranged from 32 to 55% (Table 3). Little difference was noted in values obtained by the various methods, although the authors considered the regression method to be the best estimator since it utilized geometric rather than arithmetic means. The slope of the descending right limb of the catch curve equals the instantaneous rate of mortality and can be easily converted to total annual mortality (Ricker 1975).

By year, the highest mean annual mortality rate for red porgy was 49% in 1972, and the lowest mortality was 43% in 1973. Thus, 43% of the red porgy of ages V and

Table 3.—Total annual mortality estimates for red porgy from North Carolina and South Carolina, 1972-74 (from Manooch 1975).

| Year    | Area     | Method |                    |        |                         |            | Sample size (V-oldest) | Number of age classes | Mean mortality | Mean survival |
|---------|----------|--------|--------------------|--------|-------------------------|------------|------------------------|-----------------------|----------------|---------------|
|         |          | Ricker | Robson and Chapman | Heinke | Rounsefell and Everhart | Regression |                        |                       |                |               |
| 1972-74 | Combined | 0.43   | 0.44               | 0.43   | 0.47                    | 0.44       | 9,066                  | 9                     | 0.44           | 0.55          |
| 1972    | 1        | 0.32   | 0.36               | 0.32   |                         | 0.29       | 474                    | 7                     | 0.32           | 0.68          |
|         | 2        | 0.54   | 0.55               | 0.54   |                         | 0.55       | 779                    | 7                     | 0.55           | 0.45          |
|         | 3        | 0.48   | 0.52               | 0.48   |                         | 0.55       | 1,621                  | 8                     | 0.52           | 0.48          |
| 1973    | Combined | 0.47   | 0.49               | 0.47   |                         | 0.52       | 2,883                  | 9                     | 0.49           | 0.51          |
|         | 1        | 0.38   | 0.42               | 0.38   |                         | 0.39       | 833                    | 7                     | 0.40           | 0.60          |
|         | 2        | 0.35   | 0.38               | 0.35   |                         | 0.37       | 1,299                  | 8                     | 0.37           | 0.63          |
|         | 3        | 0.44   | 0.46               | 0.44   |                         | 0.43       | 1,888                  | 8                     | 0.44           | 0.56          |
| 1974    | Combined | 0.40   | 0.42               | 0.40   |                         | 0.47       | 4,044                  | 9                     | 0.43           | 0.57          |
|         | 1        | 0.35   | 0.36               | 0.35   |                         | 0.40       | 259                    | 8                     | 0.37           | 0.63          |
|         | 2        | 0.48   | 0.47               | 0.48   |                         | 0.43       | 600                    | 7                     | 0.46           | 0.54          |
|         | 3        | 0.44   | 0.44               | 0.44   |                         | 0.42       | 1,218                  | 9                     | 0.43           | 0.57          |
|         | Combined | 0.44   | 0.43               | 0.44   |                         | 0.44       | 2,077                  | 9                     | 0.44           | 0.56          |

older were expected to die that year attributable either to harvest or natural causes.

Geographically, Cape Lookout generally had the lowest annual mortality for the 3 yr: 32%, 40%, and 37%. Overall, Cape Fear had the highest rates of total annual mortality: 55%, 37%, and 46%.

#### 4.6 The population in the community and the ecosystem

Although the habitats are fairly similar off northwest Africa, South America, and the southeastern United States, with respect to depth range, temperature, and substrate, species of fish associated with the red porgy vary. Wozinak (1967) listed the following species of sparids commonly found with *P. pagrus* off northwest Africa: *Dentex filiosus*, *D. canariensis*, *D. macrophthalmus*, *D. maroccanus*, *Chrysophrys aurata*, *Pagrus ehrenbergi*, *P. auriga*, *Pagellus canariensis*, *P. mormyrus*, *P. acarne*, *P. bogaraveo*, *Sargus vulgaris*, *S. bellotti*, *Charax puntazzo*, *Box boops*, and *Cantharus cantharus*. Klimaj (1970) also listed species of fish trawled with *P. pagrus*. His list of sparids is very similar to that above, and also includes other families and species (Table 4).

Austin (see footnote 4) mentioned the red porgy as the most commonly caught species of fish on the Florida Middle Ground, a reef located off Tampa, in the Gulf of Mexico. Physical characteristics of the area are referred to in section 2.1. An ecological condition noted by Austin was: dense carpet of brown algae during summer, disappearing in fall and winter. The only winter benthic algae observed was heavily grazed colonies of *Caulerpa pelatata*. During summer when they are abundant, herbivorous crabs such as *Mithrax pluvacantha* are major foods of demersal fish. Species of fishes found collected with red porgy are listed in Table 5.

Manooch (1975) identified concentrations of red porgy at two major depth regimes off North Carolina and South Carolina (section 2.1). Species of carnivorous fishes caught with *P. pagrus* revealed two different communities. Along the southern coast of Brazil (Espírito

Santo) large schools of *Pagrus pagrus* are found associated with dense growths of the algae *Laminaria* (Cordeiro and Miyares, pers. commun.). The species generally concentrates along mussel beds at a depth of approximately 30 m off Argentina (M. Haimovici, pers. commun., Fundação Universidade do Rio Grande).

## 5 FISHING

Since the red porgy is so widely distributed and is landed by fisheries of nations with diverse cultures in the Americas, Africa, and Europe, fishing techniques vary from one area to another. Fisheries of the southeastern United States, northwest Africa, and South America have been described and will be discussed under the headings which follow.

### 5.1 Fishing Equipment

#### 5.11 Gear

Bottom trawls, traps, and hook-line are used to capture *P. pagrus*. Off the Carolinas red porgy are taken almost exclusively by hook and line by both recreational and commercial fishermen. Huntsman (1976) described gear used aboard headboats. Tackle used on the boats is heavy and sturdy enough to resist the abuse of inexperienced anglers, and to land large fish. Fish are caught using 6/0 to 9/0 reels, either powered manually or electrically. Rods are generally constructed of fiberglass with the rod blank extending through the handle. Lines are 80- to 120-lb monofilament. Baits (squid and cut fish) are fished directly on the bottom using rigs constructed of 80-lb monofilament line with two 4/0 to 7/0 hooks. Sinkers vary from 8 to 28 oz depending on current velocity or depth. Heavy sinkers seem to increase effectiveness of bait presentation during rough seas or when currents sweep lighter weights from the bottom. "Snapper reels," which also utilize hook and line, are commonly used on snapper boats of the southeastern and gulf coasts of



Table 4.—Species of fish trawled with *Pagrus pagrus* off Mauritania and Senegal, by depth and month(s) (from Klimaj 1970).

| Family:Species                      | Depth occurred with <i>Pagrus pagrus</i> |                         |                                   |
|-------------------------------------|--|-------------------------|-----------------------------------|
|                                     | 28 Mar.-<br>18 Apr.<br>(110-135 m)       | 11-30 May<br>(40-150 m) | 20 June-<br>10 July<br>(18-120 m) |
| <b>Pomatomidae</b>                  |  |                         |                                   |
| <i>Pomatomus saltatrix</i>          | 110-135                                  | 40-150                  | 18-120                            |
| <b>Carangidae</b>                   |  |                         |                                   |
| <i>Trachurus trachurus</i>          | 110-135                                  | 40-150                  | 18-120                            |
| <i>Trachurus picturatus</i>         | 110-135                                  | 50-150                  | 18-120                            |
| <i>Decapturus punctatus</i>         | 110-135                                  | 40-150                  | 65-120                            |
| <i>Decapturus ronchus</i>           | 110-135                                  | 40-150                  | 18-120                            |
| <i>Selar crumenophthalmus</i>       | 110-135                                  | 50-150                  | 65-120                            |
| <i>Vomer setipinnis</i>             | 110-135                                  | 40-125                  | 65-120                            |
| <i>Lichia vadigo</i>                | —  | 40-150                  | 18-120                            |
| <b>Sparidae</b>                     |  |                         |                                   |
| <i>Pagrus ehrenbergi</i>            | 110-135                                  | 40-150                  | 18-120                            |
| <i>Dentex macrophthalmus</i>        | 110-135                                  | 50-150                  | 65-120                            |
| <i>Dentex filosus</i>               | 110-135                                  | 40-150                  | 65-120                            |
| <i>Pagellus centrodontus</i>        | 110-135                                  | —                       | —                                 |
| <i>Pagellus erythrinus</i>          | —  | 50-150                  | 65-120                            |
| <i>Pagellus acarne</i>              | 110-135                                  | 40-150                  | 65-120                            |
| <i>Cantharus lineatus</i>           | 120-135                                  | —                       | —                                 |
| <i>Sparus aurata</i>                | 110-135                                  | 40-150                  | 95-120                            |
| <i>Sarpa salpa</i>                  | —  | 70-150                  | 65-120                            |
| <i>Diplodus annularis</i>           | 110-135                                  | 40-80                   | —                                 |
| <i>Dentex canariensis</i>           | —  | 50-150                  | 65-120                            |
| <i>Boops boops</i>                  | —  | 40-150                  | 65-120                            |
| <i>Pagellus bogaraveo</i>           | —  | 40-150                  | 65-120                            |
| <i>Diplodus sargus</i>              | —  | 70-80                   | 65-120                            |
| <b>Sciaenidae</b>                   |  |                         |                                   |
| <i>Johnius hololepidotus</i>        | 120-135                                  | —                       | 65-120                            |
| <i>Sciaena cirrosa</i>              | 120-135                                  | 40-45                   | 65-120                            |
| <i>Sciaena ronchus</i>              | —  | 50-150                  | —                                 |
| <b>Serranidae</b>                   |  |                         |                                   |
| <i>Serranus aeneus</i>              | 120-135                                  | —                       | 65-120                            |
| <i>Serranus alexandrinus</i>        | 110-135                                  | —                       | —                                 |
| <b>Scombridae</b>                   |  |                         |                                   |
| <i>Scomber japonicus</i>            | 110-135                                  | 40-150                  | 18-75                             |
| <i>Orcynopsis unicolor</i>          | —  | —                       | 18-24                             |
| <b>Lutjanidae</b>                   |  |                         |                                   |
| <i>Brachydeuterus auritus</i>       | 120-135                                  | 70-150                  | 65-120                            |
| <i>Parapristipoma mediterraneum</i> | —  | —                       | 18-120                            |
| <b>Clupeidae</b>                    |  |                         |                                   |
| <i>Sardinella aurita</i>            | 110-135                                  | 60-125                  | 18-75                             |
| <b>Zeidae</b>                       |  |                         |                                   |
| <i>Zeus faber</i>                   | 110-135                                  | —                       | —                                 |
| <b>Gadidae</b>                      |  |                         |                                   |
| <i>Merluccius merluccius</i>        | 110-135                                  | 50-80                   | 65-120                            |
| <b>Trichiuridae</b>                 |  |                         |                                   |
| <i>Trichiurus lepturus</i>          | 110-135                                  | 40-150                  | 18-110                            |
| <b>Sphyraenidae</b>                 |  |                         |                                   |
| <i>Sphyraena barracuda</i>          | —  | 50-105                  | 18-75                             |

North America. They are usually electrically or hydraulically operated.

Apparently hand lines as well as traps have been used for many years to catch red porgy and other sparids off northwest Africa. Murray and Hjort (1912) described these techniques from the Canary Islands, and the author observed almost identical gear in 1974. Brandt (1970) referred to fishing equipment used off northwest Africa for many groups of fish including sparids: "... generally speaking the traditional fisheries in the area under discussion use lines (hand-lines and long-lines), gill nets (set nets and drift nets) and beach seines.

Traps (for fish and crustaceans), cast nets and also small surrounding nets are also well known." Trawls (bottom trawls, one-boat and two-boat systems) are also used. Detailed descriptions and illustrations of various bottom trawls as used by the Soviet Union and Poland are provided by Savrasov (1970) and Czajka and Burawa (1970), respectively. Traps, trawls, and handlines are used to capture *P. pagrus* off the east coast of South America. There are two principal Argentine fleets (Cotrina and Cousseau see footnote 3). Small trap boats fish relatively close to shore in shallow water and the catch is sold locally. Cordini (1962) provided a description of the fish traps or nasas. Larger vessels trawl for *Pagrus* in deeper waters and the fish are generally frozen and exported. Cordeiro and Miyares (pers. commun.) stated that they are collected off the southern coast of Brazil by handline. Commercial fishermen use lines that have 12 hooks.

### 5.12 Boats

Huntsman (1976) described the three basic qualities of headboats used off the southeastern United States: 1) ample passenger capacity; 2) sufficient speed to make the long trip offshore acceptable to anglers; and 3) loran and depth recording instruments to locate fish and suitable bottom topography. Passenger capacity of both inshore and offshore boats vary from 30 to 75 anglers. Speed is most essential to offshore boats and they may attain speeds up to 34.5 kn, although 23 kn is about average. Boats vary in length from 18 to 34 m.

There is a great diversity in fishing vessels used off northwest Africa. A traditional fishery with pirogues (canoes) with various modifications including sails, and row boats exist simultaneously with national and foreign fleets of big factory and freezer trawlers. Fishing vessels, then, range in size from small rowing boats to big stern trawlers and factory ships (Brandt 1970). Brandt (1970) reported that the greatest influence on fish stocks (including sparids) off the West African Shelf are the highly developed trawlers of foreign nations, especially freezer-stern trawlers of Western, Southern, and Eastern Europe as well as the Far East. Russian and Polish trawling vessels are discussed in detail by Savrasov (1970) and Czajka and Burawa (1970).

In addition to these rather large fishing vessels, there exists off northwest Africa a fleet of small line and trap boats mainly of Spanish and Mauritanian origin. South American vessels were briefly mentioned in the previous section.

### 5.2 Fishing areas

Fishing grounds off North Carolina and South Carolina are identified by Manooch (1975) and are fairly representative of the entire southeast U.S. coast: "The outer Continental Shelf of the Carolinas provides two types of habitat which serve as fishing grounds for red porgy. The most spectacular of these is the shelf break zone (Struhsaker 1969) where the ocean floor slopes



Table 5.—List of fishes observed on the Florida Middle Ground, in addition to red porgy (from Austin see text footnote 4).

| Family          | Species                          | Common name          | Family         | Species                            | Common name           |
|-----------------|----------------------------------|----------------------|----------------|------------------------------------|-----------------------|
| Caracharhinidae | <i>Carcharhinus falciformis</i>  | Silky shark          |                | <i>Decapterus</i> sp.              | Scad                  |
|                 | <i>Carcharhinus leucas</i>       | Bull shark           |                | <i>Naucrates ductor</i>            | Pilotfish             |
|                 | <i>Galeocerdo cuvieri</i>        | Tiger Shark          |                | <i>Seriola dumerili</i>            | Greater amberjack     |
| Clupeidae       | <i>Etrumeus teres</i>            | Round herring        |                | <i>Trachurus lathami</i>           | Rough scad            |
| Engraulidae     | <i>Anchoviella perfasciata</i>   | Flat anchovy         | Coryphaenidae  | <i>Coryphaena hippurus</i>         | Dolphin               |
| Synodontidae    | <i>Synodus intermedius</i>       | Sand diver           | Pomadasyidae   | <i>Haemulon aurolineatum</i>       | Tomtate               |
| Muraenidae      | <i>Gymnothorax moringa</i>       | Spotted moray        |                | <i>Haemulon plumieri</i>           | White grunt           |
| Ophichthidae    | <i>Ahlia egmontis</i>            | Key worm eel         | Sciaenidae     | <i>Equetus umbrosus</i>            | Cubbyu                |
|                 | <i>Myrophis punctatus</i>        | Speckled worm eel    | Mullidae       | <i>Mullus auratus</i>              | Red goatfish          |
|                 | <i>Ophichthus gomesi</i>         | Shrimp eel           | Sparidae       | <i>Calamus nodosus</i>             | Knobbed porgy         |
| Belonidae       | <i>Thylosurus crocodilus</i>     | Houndfish            | Ephippidae     | <i>Chaetodipterus faber</i>        | Atlantic spadefish    |
| Exocoetidae     | <i>Euleptorhamphus velox</i>     | Flying halfbeak      | Chaetodontidae | <i>Chaetodon c.f. ocellatus</i>    | Spotfin butterflyfish |
|                 | <i>Hemiramphus brasiliensis</i>  | Ballyhoo             |                | <i>Pomacanthus arcuatus</i>        | Gray angelfish        |
|                 | <i>Cypselurus heterurus</i>      | Atlantic flyingfish  |                | <i>Holocanthus bermudensis</i>     | Blue angelfish        |
|                 | <i>Parexocoetus brachypterus</i> | Sailfin flyingfish   | Pomacentridae  | <i>Chromis c.f. scotti</i>         | Purple reeffish       |
|                 | <i>Hirundichthys rondeleti</i>   | Blackwing flyingfish |                | <i>Pomacentrus c.f. variabilis</i> | Cocoa damselfish      |
| Bregmacerotidae | <i>Bregmaceros atlanticus</i>    | Antenna codlet       | Labridae       | <i>Bodianus rufus</i>              | Spanish hogfish       |
| Holocentridae   | <i>Holocentrus ascensionis</i>   | Squirrelfish         |                | <i>Halichoeres bivittatus</i>      | Slippery dick         |
| Serranidae      | <i>Centropristis ocyurus</i>     | Bank sea bass        |                | <i>Lachnolaimus maximus</i>        | Hogfish               |
|                 | <i>Diplectrum formosum</i>       | Sand perch           |                | <i>Thalassoma bifasciatum</i>      | Bluehead              |
|                 | <i>Epinephelus adscensionis</i>  | Rock hind            | Scaridae       | <i>Sparisoma</i> sp.               | Parrotfish            |
|                 | <i>Epinephelus guttatus</i>      | Red hind             | Acanthuridae   | <i>Acanthurus chirurgus</i>        | Doctorfish            |
|                 | <i>Epinephelus morio</i>         | Red grouper          | Gempylidae     | <i>Gempylus serpens</i>            | Snake mackerel        |
|                 | <i>Mycteroperca microlepis</i>   | Gag                  | Scombridae     | <i>Scomberomorus maculatus</i>     | Spanish mackerel      |
|                 | <i>Mycteroperca phenax</i>       | Scamp                | Gobiidae       | <i>Gobiosoma oceanops</i>          | Neon goby             |
|                 | <i>Petrometopon cruentatum</i>   | Graysby              | Scorpaenidae   | <i>Scorpaena plumieri</i>          | Spotted scorpionfish  |
| Lutjanidae      | <i>Lutjanus campechanus</i>      | Red snapper          | Sphyrnaeidae   | <i>Sphyrna barracuda</i>           | Great barracuda       |
|                 | <i>Lutjanus griseus</i>          | Gray snapper         | Mugilidae      | <i>Mugil cephalus</i>              | Striped mullet        |
|                 | <i>Lutjanus synagris</i>         | Lane snapper         | Bothidae       | <i>Syacium micrurum</i>            | Channel flounder      |
|                 | <i>Ocyurus chrysurus</i>         | Yellowtail snapper   | Soleidae       | <i>Gymnachirus melas</i>           | Naked sole            |
|                 | <i>Rhomboplites aurorubens</i>   | Vermilion snapper    | Cynoglossidae  | <i>Symphurus</i> sp.               | Tonguefish            |
| Priacanthidae   | <i>Pristigenys alta</i>          | Short bigeye         | Echeneidae     | <i>Echeneis naucrates</i>          | Sharksucker           |
| Apogonidae      | <i>Apogon pseudomaculatus</i>    | Twospot cardinalfish | Balistidae     | <i>Balistes capricus</i>           | Gray triggerfish      |
| Rachycentridae  | <i>Rachycentron canadum</i>      | Cobia                |                | <i>Monacanthus setifer</i>         | Pygmy filefish        |
| Carangidae      | <i>Caranx ruber</i>              | Bar jack             | Batrachoididae | <i>Opsanus pardus</i>              | Leopard toadfish      |

abruptly from the Continental Shelf to the Continental Slope. This break zone occupies an area from 30 to 100 fathoms and is characterized by very irregular topography of jagged peaks, precipitous cliffs, and rocky ledges. Less spectacular, but equally productive are the 'live bottom' areas (Struhsaker 1969) of the Continental Shelf shoreward of the shelf break area. Scattered over this submarine plain are numerous shipwrecks, rock outcroppings, and coral patches (Huntsman and Macintyre 1971) which provide ideal habitat for red porgy. Boats which fish over the latter are referred to as 'inshore' boats and those that fish the shelf break are referred to as 'off-shore' (Fig. 10) vessels."

Large quantities of *P. pagrus* and other sparids are taken commercially off northwest Africa. The following description of the major fishing area from Morocco to Senegal was taken from Brandt (1970). The area under discussion (lat. 36°N to 15°N) corresponds to the old French and Portuguese fishing areas Xa, Xb including Madeira Islands, XIa with the Canary Islands, XIb, and the northern part of XIIa, as well as Portuguese fishing area XIIc with the Cape Verde Islands to long. 28°W. This area also corresponds to the Italian fishing areas 14a, 14b with Madeira Islands, 14c with Canary Islands, and 14f. Morocco, Ifni, Rio de Oro (Spanish Sahara), Mauritania, and the northern part of western Senegal are the most important fishing areas. Coral reefs hinder bot-

tom trawling in some areas especially the northernmost coast of Morocco to Cape Juby; this area is mainly for pelagic fishing. Bottom trawling is more prevalent off




Figure 10.—Inshore and offshore fishing grounds off North Carolina and South Carolina (from Manooch 1975).




Figure 11.—Fishing grounds off Brazil (from Yesaki and Barcellos 1974).




Figure 12.—Major Argentine fishing areas (from Cotrina and Cousseau, see text footnote 3).

the coast of Rio de Oro (Spanish Sahara). This includes Arguin Bank off the northern coast of Mauritania south of Port Etienne about 18 mi off the coast. The Baie du Levrier is the nearest part of the bank from Port Etienne and one of the few sheltered areas of the coast protected by Cape Blanco. The shelf here is about 80 mi wide. The whole area from Cape Juby and the Arguin Bank is considered suitable for bottom trawling. Of the three groups of islands (Madeira, Canary, and Cape Verde) included in this large geographical area, only the Canary Islands have fisheries which are closely associated with the

northwest coast of Africa. Fishing grounds off South America are illustrated by Yesaki and Barcellos (1974) in Figure 11 and Cotrina and Cousseau (see footnote 3) in Figure 12.

### 5.3 Fishing seasons

Fishing seasons for red porgy off the southeastern United States are not regulated by law and probably not by differences in seasonal availability, but rather by weather and economic factors. Landings are not reported

on a monthly or seasonal basis for most of the fishing areas in the Atlantic. Manooch (1975) found little difference in CPUE for *P. pagrus* caught by hook and line off the Carolinas. Cotrina and Cousseau (see footnote 3) reported higher trawl catches in summer (December-March) off Argentina, but that the species is taken throughout the year by trap fishermen.

#### 5.4 Fishing operations and results

The Beaufort Laboratory, NMFS, has collected catch and effort data from North Carolina and South Carolina headboats since 1972. These data are the only species specific catch information available for red porgy landed

Table 6.—Catch of red porgy in thousands of metric tons caught off South America.<sup>1</sup>

| Year   | Country   |        |      |           | Totals |
|--------|-----------|--------|------|-----------|--------|
|        | Argentina | Brazil | USSR | Venezuela |        |
| 1938   | 0.1       |        |      |           | 0.1    |
| 1948   | 0.4       |        |      |           | 0.4    |
| 1958   | 1.6       | 0.7    |      |           | 2.3    |
| 1959   | 1.7       |        |      |           | 1.7    |
| 1960   | 1.5       |        |      |           | 1.5    |
| 1961   | 1.3       | 0.7    |      |           | 2.0    |
| 1962   | 1.4       | 0.7    |      |           | 2.1    |
| 1963   | 1.7       | 1.1    |      |           | 2.8    |
| 1964   | 2.1       | 1.2    |      |           | 3.3    |
| 1965   | 2.1       | 3.1    |      |           | 5.2    |
| 1966   | 3.1       | 3.4    |      |           | 6.5    |
| 1967   | 2.5       | 5.4    | 17.6 |           | 25.5   |
| 1968   | 2.7       | 3.8    | 11.8 | 0.3       | 18.6   |
| 1969   | 1.8       | 3.0    |      | 0.2       | 5.0    |
| 1970   | 2.1       | 3.2    |      |           | 5.3    |
| 1971   | 1.9       | 3.2    |      |           | 5.1    |
| 1972   | 1.6       | 4.7    |      |           | 6.3    |
| 1973   | 3.6       | 4.7    |      |           | 8.3    |
| 1974   | 1.4       | 4.7    |      |           | 6.1    |
| Totals | 34.6      | 43.6   | 29.4 | 0.5       | 108.1  |

<sup>1</sup>Source: FAO Fishery Statistics Yearbooks 1938, 1948, and 1958-74.

in the United States. Recreational catch data for other areas are nonexistent, and commercial landings are generally reported with several species of sparids combined. In the southeastern United States and Gulf of Mexico, *P. pagrus* is generally combined with two other genera of porgy, *Stenotomus* and *Calamus*, under the heading "scup or porgy" (Anonymous 1971). From northwest Africa *P. pagrus* is often reported under the heading "page" which includes several genera: *Pagrus*, *Sparus*, and *Pagellus* (Monteiro and Dias 1970).

#### 5.41 Effort and intensity

Off the Carolinas red porgy are caught with serranids, pomadasyids, and lutjanids, using methods described in section 5.11. Effort was reported as catch per angler day: the number and/or kilograms of red porgy caught by one angler fishing 1 day (Sekavec and Huntsman 1972; Huntsman 1976; Manooch 1975). Catch and effort data for red porgy landed by recreational fishermen off North Carolina and South Carolina are presented in Table 2. Monteiro and Dias (1970) and Zei and Ansa-Emmim (1970) provided trawling effort by the Portuguese and Ghanaian trawlers, respectively, off northwest Africa, and Yesaki and Barcellos (1974) furnished CPUE for trawlers off Brazil.

#### 5.43 Catches

As mentioned in section 5.4, *P. pagrus* is generally combined with other sparids in landings reports. Some catch data are available in FAO Fishery Statistics Yearbooks 1938, 1948, and 1958-74. These catches, by country and fishing areas, are provided in Tables 6 and 7.

### ACKNOWLEDGMENTS

We are indebted to the Sport Fishing Institute for providing financial assistance for a trip to the Canary Is-

Table 7.—Catches of red porgy in thousands of metric tons for years and areas where specific catch data are available.<sup>1</sup>

| Fishery areas and country    | Year |      |      |      |      |      |      |      | Total |
|------------------------------|------|------|------|------|------|------|------|------|-------|
|                              | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 |       |
| Atlantic, northeast          |      |      |      |      |      |      |      |      |       |
| Spain                        | 0.6  | 0.3  | 0.4  | 0.3  | 0.6  | 0.5  | 0.5  | —    | 3.2   |
| Area subtotals               | 0.6  | 0.3  | 0.4  | 0.3  | 0.6  | 0.5  | 0.5  |      | 3.2   |
| Atlantic, eastern central    |      |      |      |      |      |      |      |      |       |
| Spain                        | 4.4  | 3.8  | 2.3  | 2.1  | 1.9  | 1.6  | 1.6  | —    | 17.7  |
| Area subtotals               | 4.4  | 3.8  | 2.3  | 2.1  | 1.9  | 1.6  | 1.6  |      | 17.7  |
| Atlantic, southeast          |      |      |      |      |      |      |      |      |       |
| Angola                       | 0.5  | 0.4  | 0.7  | 1.0  | 1.2  | 0.4  | 0.4  | 0.1  | 4.7   |
| Area subtotals               | 0.5  | 0.4  | 0.7  | 1.0  | 1.2  | 0.4  | 0.4  | 0.1  | 4.7   |
| Mediterranean and Black Seas |      |      |      |      |      |      |      |      |       |
| Libyan Arab                  |      |      |      |      |      |      |      |      |       |
| Republic                     | 0.4  | 0.5  | 0.7  | 0.9  | 1.0  | 1.2  | 1.5  | 1.5  | 7.7   |
| Spain                        | 0.0  | 0.1  | 0.1  | 0.1  | 0.1  | 0.1  | 0.1  | —    | 0.6   |
| Tunisia                      | —    | —    | —    | —    | —    | 0.1  | 0.1  | 0.1  | 0.3   |
| Turkey                       | 0.4  | 0.5  | 0.3  | 0.5  | 1.1  | 0.2  | 0.1  | 0.1  | 3.2   |
| Area subtotals               | 0.8  | 1.1  | 1.1  | 1.5  | 2.2  | 1.6  | 1.8  | 1.7  | 11.8  |

<sup>1</sup>Source: FAO Fishery Statistics Yearbooks 1965-72.

lands, and the Director, C. Garcia-Cabrera, and staff of the Laboratorio Oceanográfico de Canarias, Tenerife, Canary Islands, for technical assistance, laboratory space, and living accommodations while in the Canary Islands. We thank the following scientists who provided information on the species from their respective countries: V. H. Bertullo, Instituto de Investigaciones Pesqueras, Montevideo, Uruguay; H. P. Castello, Museo Argentino de Ciencias Naturales, Buenos Aires, Argentina; S. S. Cordeiro and A. Miyares, Programa de Pesquisa e Desenvolvimento Pequeiro do Brasil, Brasilia, Brasil; C. P. Cotrina, Instituto de Biologia Marine, Mar del Plata, Argentina; N. Ehrhardt, Instituto Nacional de Pesca, Montevideo, Uruguay; J. L. Guimaraes, Ministério da Agricultura/SUDEPE, Rio de Janeiro, Brasil; M. Haimovici, Fundação Universidade do Rio Grande, Base Oceanográfico Atlântica, Rio Grande, Brasil; N. A. Menezes, Museu de Zoologia da Universidade de São Paulo, São Paulo, Brasil; G. Vazzoler, Universidade de São Paulo, Instituto Oceanográfico, São Paulo, Brasil; and M. Yesaki, SUDEPE/PDP, Rio Grande, R. S. Brasil. We are also grateful to J. R. Montalva, Puerto Rico Nuclear Center, Mayaguez, P.R., for translating manuscripts and to John W. Reintjes and James E. Sykes, National Marine Fisheries Service, Beaufort, for reviewing the synopsis.

## LITERATURE CITED

### ANONYMOUS.

1971. Fisheries statistics of the United States, 1971. U.S. Dep. Commer., NOAA, Natl. Mar. Fish. Serv., Stat. Dig. 65, 424 p.
- BEARDEN, C. M., and M. D. MCKENZIE.  
1971. An investigation of the offshore demersal fish resources of South Carolina. S.C. Wildl. Resour. Dep. Tech. Rep. No. 2, 19 p.
- BEAUMARIAGE, D. S.  
1969. Returns from the 1965 Schlitz tagging program including a cumulative analysis of previous results. Fla. Dep. Nat. Resour., Mar. Res. Lab., Tech. Ser. 59, 38 p.
- BERTALANFFY, L. von.  
1938. A quantitative theory of organic growth (inquiries on growth laws. II). Hum. Biol. 10:181-213.
- BLOCH, M. E., and J. G. SCHNEIDER.  
1801. Systema ichthyologiae iconibus CX illustratum, post obitum auctoris opus inchoatum absoluit. correxit, interpolavit. J. G. Schneider, Berlin, 572 p.
- BRANDT, A. V.  
1970. Fishing off Northwest-Africa. Rapp. P.-V. Reun. Cons. Int. Explor. Mer 159:231-239.
- CAUSEY, D.  
1953. Parasitic copepoda of Texas coastal fishes. Univ. Tex., Publ. Inst. Mar. Sci. 3:5-16.
- CIECHOMSKI, J. D., and G. WEISS.  
1973. Desove y desarrollo embrionario y larval de besugo, *Pagrus pagrus* (Linne) en el Mar Argentina (Pisces, Sparidae). Physis, Secc. A, Oceanos Org. 32:481-487.
- CORDINI, J. M.  
1962. La pesca en el Mar Argentino. Secr. Agr. Ganad. de la Nacion. Buenos Aires:1-28.
- CZAJKA, W., and W. BURAWA.  
1970. Trawl nets used by Polish trawlers on fishing grounds of the West African Shelf. Rapp. P.-V. Reun. Cons. Int. Explor. Mer 159:264-271.
- DIAS, R. K., J. K. DIAS, and W. D. ANDERSON, JR.  
1972. Relationships of lengths (standard, fork, and total) and lengths to weight in the red porgy, *Pagrus sedecim* (Perciformes,

- Sparidae), caught off South Carolina. Trans. Am. Fish. Soc. 101:503-506.
- D'ANCONA, U.  
1950. Détermination et différenciation du sexe chez les poissons. Arch. Anat. Microsc. Morphol. Exp. 39:274-294.  
1956. Inversion spontanées et expérimentales dans les gonades des téléostéens. Annee Biol. 32:89-99.
- FOWLER, H. W.  
1936. The marine fishes of West Africa based on the collection of the American Museum Congo Expedition, 1909-1915. Bull. Am. Mus. Nat. Hist. 70:607-1493.
- GINSBURG, I.  
1952. Eight new fishes from the Gulf coast of the United States with two new genera and notes on geographic distribution. J. Wash. Acad. Sci. 42:84-101.
- HILGENDORF, F.  
1888. Die Fische der Azoren. Arch. Naturgesch. 54(1):205-213.
- HUNTSMAN, G. R.  
1976. Offshore headboat fishing in North Carolina and South Carolina. Mar. Fish. Rev. 38(3):13-23.
- HUNTSMAN, G. R., and I. MACINTYRE.  
1971. Tropical coral patches in Onslow Bay. Underwater Nat. 7(2):32-34.
- JONES, P. G. W., and A. R. FOLKARD.  
1970. Chemical oceanographic observations off the coast of north-west Africa, with special reference to the process of upwelling. Rapp. P.-V. Reun. Cons. Int. Explor. Mer 159:38-60.
- JORDAN, D. S., and J. A. GUNN, JR.  
1899. List of fishes collected at the Canary Islands by Mr. O. F. Cook, with descriptions of four new species. Proc. Acad. Nat. Sci. Phila. 50:339-347.
- KLIMAJ, A.  
1970. Fishing yield and species composition of trawl catches of M. T. "Ramada" on the fishing grounds on the shelf off Mauritania and Senegal in 1967. Rapp. P.-V. Reun. Cons. Int. Explor. Mer 159:254-260.
- LINNAEUS, C.  
1758. Systema Naturae, 10th ed. Holmiae. Nantes and Pisces, p. 230-338. (Reprint, 1956, Lond.)
- MACINTYRE, I. G., and J. D. MILLIMAN.  
1970. Physiographic features on the outer shelf and upper slope, Atlantic Continental Margin, Southeastern United States. Geol. Soc. Am. Bull. 81:2577-2598.
- MANOOCH, C. S. III.  
1975. A study of the taxonomy, exploitation, life history, ecology and tagging of the red porgy, *Pagrus pagrus* Linnaeus off the Carolinas. Ph.D. Thesis, North Carolina State Univ., Raleigh, 275 p.  
1976. Reproductive cycle, fecundity, and sex ratios of the red porgy, *Pagrus pagrus* (Pisces: Sparidae) in North Carolina. Fish. Bull., U.S. 74:775-781.  
In press. Food habits of the red porgy, *Pagrus pagrus* Linnaeus (Pisces: Sparidae) off North Carolina and South Carolina U.S.A. Bull. Mar. Sci.
- MANOOCH, C. S. III, and G. HUNTSMAN.  
1977. Age, growth and mortality of the red porgy, *Pagrus pagrus*. Trans. Am. Fish. Soc. 106:26-33.
- MANOOCH, C. S. III, G. R. HUNTSMAN, B. SULLIVAN, and J. ELLIOTT.  
1976. Conspecific status of the sparid fishes *Pagrus sedecim* Ginsburg and *Pagrus pagrus* Linnaeus. Copeia 1976:678-684.
- MONTEIRO, R., and M. L. DIAS.  
1970. Sur la peche Portugaise au chalut dans la zone XI (Cote Mauritanienne) du C.I.E.M. (1955-1964). Rapp. P.-V. Reun. Cons. Int. Explor. Mer 159:240-247.
- MURRAY, J., and J. HJORT.  
1912. The depths of the ocean. Macmillan and Co., Lond., 821 p.
- RANDALL, J. E.  
1968. Caribbean reef fishes. T.F.H. Publ. Inc., Jersey City, N.J., 318 p.
- RANZI, S.  
1969. Family 3: Sparidae. In S. Lo Bianco, Eggs, larvae, and juvenile stages of Teleostei, Parts I and II, p. 330-375. Fauna and

- flora of the Bay of Naples, Monogr. 38. (Translated from Ital., TT68-50346.)
- RICKER, W. E.  
1975. Computation and interpretation of biological statistics of fish populations. Fish. Res. Board Can. Bull. 191, 382 p.
- ROSA, H., JR.  
1965. Preparation of synopsis on the biology of species of living aquatic organisms. FAO Fish. Synop. 1, rev. 1, 75 p.
- SAVRASOV, V. K.  
1970. Fishing vessels, gear and methods at the coast of Africa. Rapp. P.-V. Reun. Cons. Int. Explor. Mer 159:248-253.
- SEKAVEC, G. B., and G. R. HUNTSMAN.  
1972. Reef fishing on the Carolina Continental Shelf. Proc. 15th Annu. Int. Game Fish Res. Conf., p. 76-86.
- SOLJAN, T.  
1963. Fishes of the Adriatic (Ribe Jarrana). Fauna et Flora Adriatica, Vol. 1, Pisces. (Transl. from Serbo-Croatian, OTS, 60-21161, 428 p.)
- SPROSTON, N. G.  
1946. A synopsis of the monogetic trematodes. Trans. Zool. Soc. Lond. 25:185-600.
- STRUHSAKER, P.  
1969. Demersal fish resources: composition, distribution, and commercial potential of the Continental Shelf stocks off South-eastern United States. U.S. Fish Wildl. Serv., Fish. Ind. Res. 4:261-300.
- TRAVASSOS, H.  
1973. Pagros das aguas Brasileiras. B. Merc. Pesq., Rio de Janeiro, GB, 5:321-360.
- VALENCIENNES, A.  
1837-1844. Ichthyologie des îles Canaries, ou histoire naturelle des poissons rapportés par M. M. Webb et Berthelot. In P. B. Webb et S. Berthelot, Histoire naturelle des îles Canaries, Vol. 2, 109 p., Paris.
- WALFORD, L. A.  
1946. A new graphic method of describing the growth of animals. Biol. Bull. (Woods Hole) 90:141-147.
- WHEELER, A.  
1969. The fishes of the British Isles and north-west Europe. MacMillan and Co., Ltd., Lond., 613 p.
- WOZNIAK, S.  
1967. Rozmieszczenie ryb proznowatych (Sparidae) na szelfie Afryki ponocno - zachodniej (w okresie maj-sierpień 1963 r). Prace Morsk. [Inst. Ryb.] 14(A):7-26.
- YESAKI, M., and B. N. BARCELLOS.  
1974. Desenvolvimento da pesca do pargo - roseo ao largo da costa sul do Brasil. Programa de pesquisa e desenvolvimento pesqueiro do Brasil. PNUD/FAO. Ministerio da Agricultura/SUDEPE. Serie documentos ocasionais, no. 6, 18 p.
- ZEI, M., and M. ANSA-EMMIM.  
1970. The catch of Ghanaian long range trawlers on the shelf off Mauritania and Senegal in 1965/67. Rapp. P.-V. Reun. Cons. Int. Explor. Mer 159:261-263.



## FISHERIES SYNOPSES

This series of documents, issued by FAO, CSIRO, INP, and NMFS, contains comprehensive reviews of present knowledge on species and stocks of aquatic organisms of present or potential economic interest. The Fishery Resources and Environment Division of FAO is responsible for the overall coordination of the series. The primary purpose of this series is to make existing information readily available to fishery scientists according to a standard pattern, and by so doing also to draw attention to gaps in knowledge. It is hoped that synopses in this series will be useful to other scientists initiating investigations of the species concerned or of related ones, as a means of exchange of knowledge among those already working on the species, and as the basis for comparative study of fisheries resources. They will be brought up to date from time to time as further information becomes available.

The documents of this Series are issued under the following titles:

|       |                             | Symbol |
|-------|-----------------------------|--------|
| FAO   | Fisheries Synopsis No.      | FIR/S  |
| CSIRO | Fisheries Synopsis No.      | DFO/S  |
| INP   | Sinopsis sobre la Pesca No. | INP/S  |
| NMFS  | Fisheries Synopsis No.      | NMFS/S |

Synopses in these series are compiled according to a standard outline described in Fib/S1 Rev. 1 (1965). FAO, CSIRO, INP, and NMFS are working to secure the cooperation of other organizations and of individual scientists in drafting synopses on species about which they have knowledge, and welcome offers of help in this task. Additions and corrections to synopses already issued will also be most welcome. Comments on individual synopses and requests for information should be addressed to the coordinators and editors of the issuing organizations, and suggestions regarding the expansion or modification of the outline, to FAO:

### FAO:

Fishery Resources and Environment Division  
Aquatic Resources Survey and Evaluation Service  
Food and Agriculture Organization of the United Nations  
Via delle Terme di Caracalla  
00100 Rome, Italy

### CSIRO:

CSIRO Division of Fisheries and Oceanography  
Box 21  
Cronulla, N.S.W. 2230  
Australia

### INP:

Instituto Nacional de Pesca  
Subsecretaria de Pesca  
Secretaria de Pesca  
Secretaria de Industria y Comercio  
Carmona y Valle 101-403  
Mexico 7, D.F.

### NMFS:

Scientific Editor  
National Marine Fisheries Service, NOAA  
Auke Bay Fisheries Laboratory  
P.O. Box 155  
Auke Bay, AK 99821  
U.S.A.

Consolidated lists of species or groups covered by synopses issued to date or in preparation will be issued from time to time. Requests for copies of synopses should be addressed to the issuing organization.

The following synopses in this series have been issued since January 1975:

|          |  |                |
|----------|--|----------------|
| FIR/S111 | Synopsis of biological data on rohu, <i>Labeo rohita</i>   | June 1975      |
| FIR/S112 | Synopsis of biological data on the Norway lobster, <i>Nephrops norvegicus</i>  | September 1975 |
| INP/S2   | Sinopsis de datos biológicos sobre la tortuga golfina, <i>Lepidochelys olivacea</i>  | February 1976  |
| FIR/S115 | Synopsis of biological data on the largemouth bass, <i>Micropterus salmoides</i>   | November 1976  |
| FIR/S114 | Synopsis of biological data on scallops, <i>Chlamys (Aequipecten) opercularis</i> , <i>Argopecten irradians</i> and <i>Argopecten gibbus</i> | December 1976  |
| FIR/S113 | Synopsis of biological data on the perch, <i>Perca fluviatilis</i> and <i>flavescens</i>   | December 1977  |